



Immunological effects of Flutamide and cyproterone acetate used two Anti-prostate cancer in Rats

Shaima'a ,H.J ;

Ala'a, J.H. ;

Hussain, J.A.

University of Babylon, College of Science, Biology Department

Summary

This present study was performed to examine the effects of two drugs of anticancer of prostate (flutamide & cyproterone acetate) on some immunological in albino male rats. The animals were divided into two groups, first treatment groups gives orally flutamide at 25 and 15 mg/kg/day and two group gives orally cyproterone acetate at 10 and 5 mg/kg/day, and control divided into two groups, first group gives normal saline and two group gives corn oil and all groups gives orally for 28 days. study immunological parameters including the concentration IgG, IgM, IgA, γ -globulin, IL-4, IL10 and measuring phagocytic activity also measuring total protein. Furthermore, The effects of these drugs have been also tested on some blood components including; WBC, differential WBCs in treatments and control groups (normal saline & corn oil) of rats.

The flutamide (25,15 mg/kg) and cyproterone acetate (10,5 mg/kg) produced significant decreases in total protein except WBC count which show significantly increases, but both drugs above don't effect in, IgA, IgM, IL-10 and phagocytic activity, on the other hand Flu (25 mg/kg) only produced significantly increase in level of IL-4 and significantly decrease in IgG and globulin- γ concentration, furthermore, CPA (10 mg/kg) cause only significantly increase in concentration of IgG and neutrophils and significantly decrease in lymphocytes. The results of this study refer that the two drugs show significant change in blood components in treated rats compared with control groups, high dose of Flu showed significant change in immunological parameters.

Introduction

The prostate cancer has been one of the most common disease for the men in United State of America after pulmonary and colon cancers (Jemal *et al.*, 2006). In Europe, this disease is more prevalence than pulmonary cancer (Boyle and Ferlag, 2005). There are many methods for the therapy of prostate cancer: 1. Androgen deprivation therapy by using hormones, and this method was used for treatment of non-transitional diseases in especially after radical therapy or prostatectomy (Aus *et al.*, 2005).

2. Androgen antagonist therapy, which binding to androgen receptor and therefore, block the androgen (Petrylak, 2002).

3. Testectomy by surgery but the side effects of it was losing of libido (Iversen *et al.*, 2000).

The flutamide and cyproterone acetate drugs represented steroid and steroid antagonists respectively, and both of them acts competitor for androgen receptor, but steroid antagonist also has progesterone activities that minimum the level of testosterone which lead to prevent the erection and other sexual performance (Anderson, 2003). Steroid antagonist dose not decrease testosterone level, therefore, the sexual performance will remain normal (Iversen *et al.*, 2001; Migliari *et al.*,



1999). Using androgen antagonist therapy is increase due to their effects and acceptance from paients compared with surgery in prostate cancer therapy therefore, they represnts an alternative therapy for maintaining of sexual functions in patients (McLeod *et al.*, 2006).The chemotherapy has aside effects not only on endocrine and reproductive systems, but also on the immune system (Crisp *et al.*, 1997). Chryssikopoutos *et al.*, (1997) reported that acomplex relationships between the immune system and glandular nervous system. The immune cells secrete several hormonal peptides and protein, such as growth hormone, lactogenic hormone, thyroid stimulating hormone, and adrenal corticotrophin hormone. Some of these hormones considered as suppressors or stimulators for the immune responses (Gaillard, 1995).

Certain cytokines are produced by immune cells which have the ability to alter certain functions of glandular nervous system (Rivest and Laflamme, 1995). In recent years, there is an increase in using flutamide by adolescent for treatment acne and hairsitism treatments (De Amorim *et al.*, 2005). The main aim of the present study is to determine the side effects of the widely of used two drugs, flutamide and cyproterone acetate on the immune system of rats by measuring the following:

1-Estimation the concentration of immunoglobulin (IgA, IgG, IgM) and anti-inflammatory cytokines (IL-4, IL-10).

2-Estimation the concentration of total protein and γ -globulin.

3-Evaluation the phagocytic activity.

Materials and Methods

1-Laboratory Animals:

Thirty mature native rats *Rattus rattus* have been brought from animal house of Univeristy of Kufa. Their blood were tested for bacterial pathogens and their respective specific antibodies , then rats with negative results were kept through out experimentation periods at lididum for ration and housing (Schneider *et al.*, 1990). The average weight of such animals ranged between 300-350 gm and their ages ranged from 4-5 mouths.

2-Preparation of Drug:

The cyproterone acetate was obtained from pharmacy and their equipment from company of Schering AG Germany/Allemague as tablets in concentration 50 mg/kg, while the equipment of flutamide drug from company Schering plough Lab.N.N. Belgium as tablets in concentration 250 mg/kg. The tablets were macerated by blender and dissolved by absolute alchol, then kept for dryness, after that corn oil was added as well as mixed and the concentration for experiments were done according to the doses for human (Sanchez-Criado *et al.*, 1990).

3- Immunization protocol:

The rats were divided into three groups , 10 replicates for each. The first treated group was divided in two subgroups , the first and second were orally received flutamide at concentration 15 and 25 mg/kg/day respectively for 28 day , while the second treated group were orally received cyproterone acetate at concentration 5 and 10 mg/kg/day for first and second subgroups consuctively and also for 28 day . The



last group (control) was divided also into two subgroup, the first and second were orally normal saline (negative control) and corn oil(positive control) respectively for 28 day.

4- Blood samples:

The blood samples were collected directly from the treated and controlled rats by heart puncture, seven day after the last immunization. The small amount of these samples were kept in sterile tube containing anticoagulant (heparin) AFM-Dispo and placed in refrigerator at 4 C° in order to measure the phagocytic activity. The remainder of these samples were kept in sterile centrifuge tubes (without anticoagulant) to separate the sera, and then measuring the concentration of the total protein, γ -globulin, immunoglobulins (IgA, IgG, IgM) and anti-inflammatory cytokines (IL-4, IL-10).

5-Immunological Assay:

5-1 Determination of serum protein :

The concentration of total protein (g/l) was estimated according to manual procedure of linear chemical procedure (company of Almacen Joaquim . Costa, Motagat, Barcelone, Spain), while the concentration of γ -globulin was evaluated according to Bishop *et al.*, (1985).

5-2 Single Radial Immunodiffusion Assay:

This test was used to determine the concentration of IgM, IgG and IgA according to Mancini *et al.*, (1965).

5-3 Determination of IL-4 and IL-10:

The concentration of IL-4 and IL-10 were determined by ELISA according to kit procedure (Company of Endogen, Canada).

5-4 Phagocytic activity test:

The means of phagocytosis by phagocytes for *Candida albicans* were measured according to Metcalf *et al.*, (1986) with certain modifications. The test was done by mixed fresh blood of immunized and controlled rats, with suitable amount of *C. albicans* isolate, then incubated at 37 C°, after that prepare the slide, stain and examine under oil immersion microscope. The percentage of phagocytic activity were calculated according to the following equation:

$$= \frac{\text{number of phagocytes for } C. \text{albicans}}{\text{Total number}} \times 100$$

6- Biometry:

The results for experiments were analyzed using statistical (programme spss version 17, and the mean, standard error as well as significant differences were done by using one-way anova (Joda, 2008).



Results

There were a decreased in mean of concentration of total protein in treated groups in comparison with control groups , and there is a significant differences at ($P<0.05$) compared with control groups , also the means of concentration of γ -globulin in animals primed by drugs were lower then that in control groups, but the concentration in rats immunized by cyproterone acetate 10 mg/kg were higher in comparison with other treated groups and there is a signhfcant differences between animals primed with flutamide 25 mg/kg and control groups.Meantime, there were no significant differences at ($P<0.05$)in percentage of phagocytic activity between treated and controlled groups (Table 1).

Table (1) : the concentration of total protein (gm/dl) and γ -globulin (gm/dl) as well as the percentages of phagocytic activity in rats primed orally by flutamide and cyproterone acetate drugs.

parameter Nature of treatment	Total protein gm/kg Mean $\pm$ S.E.	γ -globulin gm/kg Mean $\pm$ S.E.	Percentage of phagocytic activity %Mean $\pm$ S.E.	probability
Negative control (normal saline)	7.25 $\pm$ 0.56	3.73 $\pm$ 0.29	38.39 $\pm$ 1.16	—
Positive control (corn oil)	6.55 $\pm$ 0.38	3.43 $\pm$ 0.56	40.03 $\pm$ 2.81	—
Flutamide 25mg/kg	*5.43 $\pm$ 0.24 **	*2.09 $\pm$ 0.05 **	42.23 $\pm$ 3.76	0.05
Flutamide 15mg/kg	*5.66 $\pm$ 0.47 **	2.75 $\pm$ 0.22	35.58 $\pm$ 2.38	0.05
Cyproterone acetae 10mg/kg	*5.79 $\pm$ 0.32 **	3.06 $\pm$ 0.37	37.87 $\pm$ 2.9	0.05
Cyproterone acetate 5 mg/kg	*5.99 $\pm$ 0.04 **	2.86 $\pm$ 0.32	42 $\pm$ 5.33	0.05

*: significant difference in comparison with negative control.

** : significant difference in comparison with positive control.

The animals treated with flutamide reveled decreased at significant ($P<0.05$) in concentration of IgG (mg/dl) in comparison with control groups , while rats treated with cyproterone acetate 10 mg/kg showed increases at significant ($P<0.05$) compared



with untreated groups . The concentration of IgM (mg/dl) showed no significant differences at ($P<0.05$) between treated and control groups , but their was decreased in concentration of this antibody in animals primed with cyproterone acetate drug, futheremore, there were no significant differences at ($P<0.05$) in concentration of IgA (mg/dl) between immunized and control groups as illustrated in table (2).

Table (2): The concentration of immunoglobulins IgM, IgG and IgA in rats primed orally by flutamide and cyproterone acetate.

parameter Nature of treatment	Concentration of IgM mg/dl Mean±S.E.	Concentration of IgG mg/dl Mean±S.E.	Concentration of IgA mg/dl Mean±S.E.	probability
Negative control (normal saline)	94.63±8.99	1714.26±175.38	238±58.99	—
Positive control (corn oil)	93.03±9.44	1810.86±144.64	265.03±27.70	—
Flutamide 25mg/kg	107.3±16.71	*810.40±42.20 **	223.53±75.53	0.05
Flutamide 15mg/kg	100.4±17.22	*1021.10±248.31 **	234.43±25.84	0.05
Cyproterone acetate 10mg/kg	73.9±12.19	*2432.20±95.80 **	184.33±106.08	0.05
Cyproterone acetate 5 mg/kg	82.73±19.55	1726±285.07	263.13±67.71	0.05

*: significant difference in comparison with negative control.

** : significant difference in comparison with positive control.

Table (3) illustrated increases in concentration of IL-4 pg/ml in treated group by flutamide 25 mg/kg in comparison with control groups, as well as there were no significant differences at ($P<0.05$), but there were no significant differences at ($P<0.05$) between other treated groups and untreated groups. Meantime , the means of IL-10 pg/ml reveled no significant differences between primed and control groups.

Table (3) : The concentration of IL-4 and IL-10 pg/ml in rats immunized orally by flutamide and cyproterone acetate drugs.

parameter Nature of treatment	Concentration of IL-4 pg/ml	Concentration of IL-10 pg/ml	Probability
----------------------------------	--------------------------------	---------------------------------	-------------



	Mean±S.E.	Mean±S.E.	
Negative control (normal saline)	49.31±1.28	27.79±8.15	—
Positive control (corn oil)	49.64±1.82	24.47±13.99	—
Flutamide 25mg/kg	*73.76±7.24 **	20.49±3.04	0.05
Flutamide 15mg/kg	50.8±5.11	23.81±6.33	0.05
Cyproterone acetate 10mg/kg	60.79±6.24	22.28±8.44	0.05
Cyproterone acetate 5 mg/kg	61.27±3.6	27.78±4.78	0.05

*: significant difference in comparison with negative control.

** : significant difference in comparison with positive control.

Discussion

The means of concentration of total protein show a decreases in treatments compared with control groups at the same time the concentration of γ - globulin is lower in treatments compared with control groups (table 1). Duques de Amorim *et al.*, (2005) fond that the total protein and γ -globulin was natural in patients used flutamide for treatment of acne, also the decreases in level albumin , which is the major protein in serum probably related with lower concentration of total protein. The decrease in means of γ -globulin may be associated with level of IgG, which represents the major component of γ -globulin .However, the concentratins of cyproterone acetate 2 and 5 mg/kg don't inhibited the synthesis of globulin- α_2 (Roy, 1976). There is no significant differences at ($P<0.05$) in phagocytic activity between treatments and control groups. Srinivasan *et al.*, (1997) reported that the flutamide alone was unstimulated for polymorphounuclear cells, but in low concentration increases release the myeloperoxidase, in addition, high concentration of this drug inhibits the activation of phagocytes in presence of phorbol myristate (PMA), also both drugs donot effect on expression Fc γ R on surfaces of phagocytes (Gomez *et al.*, 2000), and this receptor mediated the phagocytosis by opsonization of molecules as well as stimulate this cells to degrade engolved particles (Abbas *et al.*, 2003).

Table (2) show a significant increase and decrease in levels of IgG in rats primed by cyproterone acetate and flutamide respectively, while there is no significant difference in levels of IgM and IgA. The low concentration of IgG in animals primed by flutamide may be due to the increase production of IL-4, which inhibits the synthesis of IgG₂ and IgG₃ (Abbas *et al.*, 2003), futhermore, the high level of IgG



related with increases activity of gamma glutamyl transferases (GGT) which elevated the amount of intracellular glutathione that lead to stimulate T-cells and leukocytes, meantime, the amount of this enzyme increased by using cyproterone acetate (Gonzalez-Quintela *et al.*, 2003). The intraperitoneal inject of rats by flutamide at 5 and 20 mg/kg for 15 days wasn't effect on level of IgM (Ladics *et al.*, 1998). Q'conner *et al.*, (2002) illustrated that the exposure of rats for flutamide 5, 20, 50 mg/kg was not effect on primary immune response . There is a significant increases in level of IgM in rats primed by cyproterone acetate at 25 mg/kg , and this is may be due to uses high doses of drugs (Ali *et al.*, 2008). Giltay *et al.*, (2000) found that decreases in level of IgA by using CPA drug at 100 mg/kg for three months in patients with sexopathy and interpretation for this case was increased the water in the body as well as plasma which lead to decrease in level of IgA, while Sande and Comp (1983) reported that increases in concentration of IgA in prostate tissue of patients with prostate cancer used CPA, because the immunoglobulin IgA was higher in mucosal tissues than peripheral blood. Table (3) showed increases in level of IL-4 in rats primed by flutamide 25 mg/kg compared with control groups while there is no significant difference in level of IL-10 between immunized and control groups. The increases in level of IL-4 may be related to elevate concentration of IL-2 which stimulate other T-cells to produce cytokines such as IFN- γ and IL-4 (Abbas *et al.*, 2003), also mice primed by flutamide at 25 mg/kg increases production of IL-2, and the later stimulate increases level of IL-4 (Messingham *et al.*, 2001; Wichmann *et al.*, 1997). Vo *et al.*, (2009) showed that rats immunize by flutamide at 10 and 50 mg/kg increased the expression of IL-4 receptor, while the levels of IL-10 was not effected in mice prime by flutamide at 25 mg/kg (Sønder *et al.*, 2006; Hildebrand *et al.*, 2006), futhermore the decreases level of IL-10 in mice showed suppression the immune response for intestinal antigen, due to binding of IL-10 to their receptors on T-cells surfaces suppress the gene transcription of IL-2 which inhibits the proliferation of T-cells (Delves and Roitt, 1998). Testosterone was revealed anti-inflammatory effect by stimulate production of IL-10 as well as inhibited production of TNF- α (D'Agostino *et al.*, 1999), therefore, the antagonist for androgen may be suppress the level of IL-10. The significant of this paper lies in the investigation of the humoral and cellular immune response to two anti-prostate cancer in rats that mimics their actions in human.

References

- Abbas, A.; Lichtman, A. & Pober, J. (2003): Cellular and Molecular Immunology. 5th ed Philadelphia., W.B. Saunders Company London.
- Ali, E.H.A. (2008): Protective effects of *Echinacea* on cyproterone acetate induced liver damage in male rats. Pakistan J. Biol. Sci., 11(21):2464-2471.
- Aus, G.; Abbou, C.C.; Bolla, M.; Heidenreich, A.; Van Poppel, H.; Schmidt, H-P.; Wolff, J.M. & Zattoni, F. (2005): Association of Urol., guidelines on prostate cancer (Euro).
- Bishop, M.G.; Dben-Von Laufer, J.L.; Fody, E.P. & Thirty three contributors (1985): Clinical Chemistry Principles, Produres and Correlations. The Murray Printing Company. Philadelphia., P: 181-182.
- Boyle, P. & Ferlay, J. (2005): Cancer incidence and mortality in Euro, 2004. Ann. Oncol., 16:481-488.



- Chryssikopoulos A** (1997): The relationship between the immune system and endocrine systems. In *Adolescent Gynecology and Endocrinology: Basic and Clinical Aspects* (G. Creatsas, G. Mastorakos, GP Chrousos, Eds.). 83–93. New York Academy of Sciences, New York.
- Crisp, T.M.; Clegg, E.D. & Cooper, R.L.** (1997): Special report on environmental endocrine disruption: An effects assessment and analysis. prepared for the risk assessment forum, U.S. Environmental Protection Agency, Washington, DC. EPA/630/R-96/012 February 1997. U.S. E.P.A .,Washington D.C.
- D'Agostino, P.; Milano, S.; Barbera, C.; Di Bella, G.; La Rosa, M.; Ferlazzo, V.; Farruggio, R.; Miceli, D.M.; Miele, M.; Castagnetta, L. & Cillari, E.** (1999): Sex hormones modulate inflammatory mediators produced by macrophages. *Ann. N.Y. Acad. Sci.*, 876:426-429.
- De Amorim, M.F.D.; Dias, A.; WaldirPped rosa; Duques, P.; Duques, D.P. & Ramalho de Vasconcelos, J.** (2005): Flutamide induced hepatotoxicity during treatment of acne-Acase report. *An. Bras. Dermatol.*, 80(4): 381-4.
- Delves, P.J. & Roitt, I.M.** (1998): *Encyclopedia of Immunology*. 2nd Ed. San Diego., Academic Press.
- Duques de Amorim, M.; Dias de Amorim, W.P.; Duques, P.; Duques de Amorim, P. & Ramalho de Vasconcelos, J.** (2005): Flutamide- induced hepatotoxicity of acne . *An. Bras. Dermatol.*, 80: 571-8.
- Gaillard, R.C.** (1995): Interactions immunoendocriniennes au niveau hypothalamo-hypophysaire. Immuno-endocrine interactions at the hypothalamo-hypophyseal level. *Paris Ann. Endocrinol.*, 56: 561–566.
- Giltay, E.J.; Frank, J.C.; Von Blombery, B.M.; Drexhage, H.A.; Schalk Wijk, C. & Gooren, L.J.** (2000): In vivo effects of sex steroids on lymphocyte responsiveness and immunoglobulin levels in humans. *J. Clin. Endocrinol. Metab.*, 85(4):1648-57.
- Gomez, F.; Ruiz, P.; Lopez, R.; Rivera, C.; Romero, S. & Bernal, J.A.** (2000): Effects of androgen treatment on expression of macrophage Fcγ receptors. *Clin. Diagn. Lab. Immunol.*, 7:682-686.
- Gonzalez-Quintela, A.; Alende, R.G. & Gonzalez-Gill, P.** (2003): Serum immunoglobulins (IgG, IgA, IgM) in chronic hepatitis C, A comparison with non-cirrhotic alcoholic liver disease. *Hepato. Gastroenterol.*, 50: 2121-2126.
- Hildebrand, F.; Thobe, B.M.; Hubbard, W.J.; Choudhry, M.A.; Pape, H.C. & Chaudry, I.H.** (2006): Effects of 17β-estradiol and flutamide on splenic macrophages and splenocytes after trauma-hemorrhage. *J. Cytokine.*, 36:107-114.
- Iversen, P.; Melezinek, I. & Schmidt, A.** (2001): Nonsteroidal antiandrogen : a therapeutic option for patients with advanced prostate cancer who wish to retain sexual interest and function . *B.J.U. Int.*, 87:47-56.



- Iversen, P.; Tyrrell, C.J.; Kaisary, A.V.; Anderson, J.B.; Van Poppel, H.; T-Ammela, T.L.; Chamberlain, M.; Carroll, K. & Melezinek, I.** (2000): Bicalutamide monotherapy compared with castration in patients with nonmetastatic locally advanced prostate cancer: 6.3 years of follow up. *J. Urol.*, 164:1579-1582.
- Jemal, A.; Siegel, R.; Ward, E.; Murray, T.; Xu, J.; Smigal, C. & Thun, M.J.** (2006): Cancer statistics, 2006. *C.A. Cancer J. Clin.*, 56:106-130.
- Joda, M.** (2008): Statistical analysis spss. Amman for publishing and distribution.
- Ladics, G.C.; Smith, C.; Nicastro, S.C.; Loveless, S.E.; Cook, J.C. & O'conner, J.C.** (1998): Evaluation of the primary humoral immune response following exposure of male rats to 17- β estradiol or flutamide for 15 days. *Toxicol. Sci.*, 14: 75-82.
- Mancini, G.; Carbonaro, A.O. & Heremans, J.F.** (1965): Immunochemical quantitation of antigens by single radial immunodiffusion. *Immunochemistry.* 2:235-54.
- McLeod, D.G.; Iversen, P.; See W.A.; Morris, T.; Armstrong, J.; Wirth, M.P.; on behaaalf of the "Casodex" Early Prostate Cancer Trialis T.S. & Grop.** (2006): Bicalutamide 150 mg plus standard care versus standard care alone for early prostate cancer . *B.J.U. Int.*, 97:247-254.
- Messingham, K.A.N.; Shirazi, M.; Duffner, L.A.; Emanuele, M.A. & Kovacs, E.J.** (2001): NT testosterone receptor blockage restores cell-ular immunity in male mice after burn injury . *J. Endocrinol.*, 169: 299-308.
- Metcalf, J.A.; Gallin, M.D.; Nauseef, M.D. & Root, R.K.** (1986) Laboratory Manual of Meutrophil Function. Raven Press ., Newyork U.S.A .
- Migliari, R.L Muscas, G.; Murru, M.; Verdacchi, T.; De Benedetto, G. & De Angelis, M.** (1999): Antiandrogen :a summary review of pharmacodynamic properties and torlerability in prostate cancer therapy .*Arch. Ital. Urol. Androl.*, 71:293-302.
- O'conner, J.C.; Frame, S.R. & Ladics, G.S.** (2002): Evaluation of a 15-day screening assay using intact male rats for identifying antiandrogen. *Toxiol. Sci.*, 69: 93-116.
- Petrylak, D.P.** (2002): Chemotherapy for the treatment hormone- refractory prostate cancer. *J. Euro. Urol. Supp.*, 2:15-23.
- Rivest, S. & Laflamme, N.** (1995): Neuronal activity and neuropeptide gene transcription in the brains of immune-challenged rats. *J. Neuroendocrinol.*, 7:501-525.
- Roitt, I.; Brostoff, J. & Male, D.** (2000): Immunology., World Moscow 63-64, 168-193.
- Roy, A.K.** (1976): Effect of cyproterone acetate on androgen- dependent synthesis of alpha 2 globulin in rats. *J. Endocrinol.*, 70: 189-95



- Sanchez-Criado, J.E.; Bellido, C.; Tebar, M.; Ruiz, A. & Gonzalez, (1999).** The antiprogesterin Ru 486 dissociates LH and FSH secretion in male rats: evidence for direct action at the pituitary level. *J. Endocrinology*, 160:197-203.
- Sande, M.V. & Comp, K.V. (1983):** Immunoglobulins in tissue of humans benign prostatic hypertrophy. *Urol. Res.* 11:171-172.
- Sønder, S.U.S.; Mikkelsen, M.; Rieneck, K.; Hedegaard, C.J. & Klaus, B. (2006):** Effects of spironolactone on human blood mononuclear cells: mineralocorticoid independent effects on gene expression and late apoptosis induction. *Br. J. Pharmacol.*, 148(1):46-53.
- Srinivasan, R.; Buchweitz, J.P. & Ganey, P.E. (1997):** Alternation by flutamide of neutrophil response to stimulation implications for tissue injury. *Biochem. Pharmacol.*, 53:1179-1185.
- Schneider, E.; Volecker, F. & Hsude, W. (1990):** Age and sex dependence on phospholipids concentration in human erythrocytes. *I.H.Med. Lab. Diagan.*, 31: 86-89.
- Vo, T.T.; Jung, E.M.; Dang, V.H.; Jung, K.; Baek, J.; Chol, K.C. & Jeung, E.B. (2009):** Differential effects flutamide and Di (2-ethylhexyl) phthalate on male reproductive organs in a rat model. *J. Rep. Dev.*, 55(4):400-411.
- Wichmann, M.W.; Angele, M.K.; Ayala, A.; Cioffi, W.G. & Chaudry, I.H. (1997):** Flutamide: A novel agent for restoring the depressed cell-mediated following soft-tissue trauma-hemorrhage. *J. Shock.*, 8:242-248.

التأثير المناعي للفلوتاميد و خلايا السايبروتيرون المضادين لسرطان الموثة (البروستات) في الجرذان

شيماء حسين جابر د. علاء جواد حسن د. حسين جاسم الحربي

الخلاصة

صممت الدراسة الحالية لبحث فعالية عقارين مضادين لسرطان الموثة (الفلوتاميد FLU و خلايا السايبروتيرون CPA) على بعض المعايير المناعية في ذكور الجرذان البيض. قسمت الحيوانات الى مجموعتين المجموعة الاولى جرعت بالفلوتاميد بالتركيزين 25 و 15 ملغم/كغم/يوم والمجموعة الثانية جرعت بخلايا السايبروتيرون بالتركيزين 10 و 5 ملغم/كغم/يوم، اما السيطرة قسمت ايضا الى مجموعتين الاولى جرعت بالمحلول الفسلجي الملحي والثانية بزيت الذرة وكل المجاميع جرعت لمدة 28 يوم. حيث درست المعايير المناعية المتمثلة بتركيز الكلوبولينات المناعية IgG, IgA, IgM وكاما كلوبيولين والسايتركائينات IL-4, IL-10 والعد التفرقي لخلايا الدم البيض والفعالية البلعمية في دم الجرذان السليمة والمجوعة بتركيز العقارين وكذلك الجرعة بزيت الذرة. لقد تبين أن عقار الفلوتاميد بتركيزي (15, 25 ملغم/كغم/يوم) وعقار خلايا السايبروتيرون بتركيزي (5, 10 ملغم/كغم/يوم) أظهر انخفاضا معنويا في البروتينات الكلية بينما أظهر ارتفاعا معنويا في عدد خلايا الدم البيض الكلي، اما عند حقن الحيوانات FLU (15, 25 ملغم/كغم) لم يؤثر معنويا على IL-10, IgM, IgA. كما ان FLU (25 ملغم/كغم) فقط أدى الى ارتفاع معنوي في IL-4 وانخفاض معنوي في IgG وكاما كلوبيولين وكذلك فقط CPA (10 ملغم/كغم) أظهر ارتفاع معنوي في IgG والخلايا متعددة الانوية وانخفاض معنوي في الخلايا اللمفاوية. يستنتج من خلال نتائج الدراسة ان عقاري FLU و CPA لهما تأثيرات سلبية على المناعة في الحيوانات المجوعة مقارنة بحيوانات السيطرة، بينما التركيز العالي من FLU له تأثيرات مناعية أكثر.