



Molecular detection of some virulence gene associated with pathogenicity of *Klebsiella pneumoniae*

*Dr. Lamees Abdul-Razzak A. Lateef
Mohammed Sabri Abdul-Razzaq

Lecturer

*Dr. Maysa Salih Mahdi

Lecturer

*Dr.

Professor

*Babylon University/ College of Medicine

Abstract:-

In this study, only seven isolates of *Klebsiella pneumoniae* were subjected for detection espB, cesT/eae, and CNF1 genes in their genome. It was seen that esp gene was not found in all isolates under study, whereas cytotoxic necrotizing factor-1 (cnf1) was detected in four isolates of *Klebsiella pneumoniae*. Moreover, three isolates were shown to possess cesT/eae gene in their genome. The presence of such genes will provide the bacteria to invade and evade the host immune system and also will interfere with wound healing as a result of cnf1 production.

Introduction:-

Many virulence genes have been sequenced in enterobacteriaceae to promote their pathogenicity such as espB, cesT/eae, and cytotoxic necrotizing factor-1 (CNF1).

The virulence determinants required for the induction of the attaching-effacing (A/E) lesion in enteropathogenic *E. coli* are encoded in a 35.6-kb pathogenicity island, denoted LEE (for locus of enterocyte effacement) (1). LEE-encoded genes have been divided into three functional regions: the *esc* (extended-spectrum cephalosporin) and *sep* genes, which code for a type III secretion-translocation system (2); the *tir* (translocated intimin receptor), *cesT* (chaperon for type III secretion of *tir*), and *eae* genes (intimate attachment to epithelial cell), coding for the proteins involved in intimate attachment (3); and the *esp* genes (intimate secreted protein), which encode effector proteins that are involved in the formation of a translocon for delivering effector molecules to the host cell (4). The *cesT* gene may encode a chaperone protein and is highly conserved in all the A-E positive strains, probably because it contains the transcriptional start of *eae* (5). On the other hand, *cesT*, previously known as *orfU*, encodes for a chaperone that is required for the stable secretion of Tir (translocated intimin receptor), it was demonstrated that *tir*, *cesT*, and *eae* constitute an operon and that *orf19* is a monocistronic unit (1). The *espB* gene has been sequenced in several strains, including human enteropathogenic *E. coli* (EPEC) (6), human and bovine enterohaemorrhagic *E. coli* (EHEC) (7), and rabbit EPEC (8). Comparison of the deduced amino acid sequences has revealed that, like intimins, EspB are highly variable proteins. The variability in genes encoding proteins that interact directly with the host, in contrast to other LEE encoded proteins, which are highly conserved, suggests that these variable proteins are subject to selection for evasion of the host immune system (9). EspB protein is essential for signal transduction in host cells and A/E lesion formation (10). However its function has not been elucidated. Mutations in the *espB* gene eliminate the formation of the A-E lesion (11).

Besides CNF-1 is a toxin produced by uropathogenic *E. coli* strains that mediates its effects via the activation of the Rho GTPases. CNF-1 can either hinder cells from apoptosis or even induce apoptosis depending on the cell type and the dose. Moreover, by influencing the phagocytosis and adherence of polymorphonuclear leukocytes, CNF-1 could facilitate the growth of bacteria in the urinary tract. In



addition, CNF-1 can enhance the secretion of proinflammatory cytokines in human uroepithelial cells. Despite many in vitro effects, there have been conflicting results about the role of CNF-1 in vivo (12).

In a study conducted by (13) who found CNF-1 was able to interfere with wound healing in intestinal epithelial monolayers.

After mechanical injury, it was found that CNF-1 blocks epithelial wound repair within 48 hours (14). This effect was characterized by cell elongation and filopodium formation on leading edge, in association with permanent phosphorylation of focal adhesion kinase via Rho activation (15). In addition to the promotion cell spreading, cnf1 protects cell from experimentally induced rounding up and detachment and improves the ability of cells to adhere to each other and to extracellular matrix by modulating the expression of proteins related to cell adhesion (16).

This study was carried out to detect the presence of *espB*, *cesT/eae*, and *CNF1* genes in local isolates of *Klebsiella pneumoniae*.

Materials and methods:-

Patients:

A total 65 samples, only seven isolates of *Klebsiella pneumoniae* were obtained from patient with burn wound infection by standard bacteriological methods. All samples were obtained from patients or individuals who admitted to Al-Hilla Surgical Teaching Hospital in Babylon city during the period from 10/2011 to 1/2012.

Bacterial identification:

The samples were processed on MacConkey agar and were incubated at 37°C overnight. The identification of gram negative bacteria was performed by standard biochemical methods (catalase test, oxidase test, and IMViC test, capsule test) according to Bergy's Manual for Determinative Bacteriology (17).

DNA extraction for gram negative bacteria:

DNA extraction was carried out according to the genomic DNA purification kit supplemented by manufactured company (promega, USA) (cat# A 1120) with modification in step (3), which involved putting in temperature (100 °C) for 10 min instead of 80 °C for 5 min.

Detection of some pathogenicity islands markers by PCR:

The primers and PCR conditions used to amplify genes encoding virulence factors with PCR are listed in table (1). The primers for the LEE include *cesT/eae* gene and *espB*, as well as the primer specific for the CNF1. Each 25 µl of PCR reaction contained 2.5 µl of each upstream and downstream primer, 2.5 µl of free nucleic acid water, 5 µl of DNA extraction and 12.5 µl of master mix. The PCR amplification product were visualized by electrophoresis on 1% agarose gels for 45 min at 60 V. The size of the amplicons were determined by comparison to the 100 bp allelic ladder (promega, USA) (cat# G 2101).

**Table (1):- Primers sequences and PCR condition to detect LEE & CNF1**


Genes	Primer sequence (5'-3')	Size of product bp	PCR condition	Reference
<i>cesT/eae</i> F <i>cesT/eae</i> R	GTTTGCAGAGAATGGTGGCCC TAGCTTATGCTTGTGCCGGGT	333	94°C 4min 1x	1
			90°C 1min 56°C 1min 30x	
			72°C 1min	
			72°C 5min 1x	
<i>espB</i> F <i>espB</i> R	GCCGCTCTGATTGGTGGTGCT TGGCGTTGAACCGGAAATCCT	387	94°C 4min 1x	1
			94°C 1min 56°C 1min 30x	
			72°C 1min	
			72°C 5min 1x	
<i>CNF1</i> F <i>CNF1</i> R	AAGATGGAGTTTCCTATGCAAGGAG CATTGAGATCCTGCCCTCATTATT	410	94°C 4min 1x	18
			94°C 1min	
			60°C 1min	
			30x	
			72°C 1min	
			72°C 7min 1x	

Company of primers is alpha (USA).

Results and Discussion:-

A total of 65 wound swabs, only seven isolates of *Klebsiella pneumonia* were identified, according to the cultural, biochemical behavior. These seven isolated were subjected to study the presence of three important virulence genes; include *espB*, *cesT/eae*, and *CNF1* genes by using PCR technique.

it was shown that the *espB* gene was not detected in all *Klebsiella pneumoniae* isolates. This may indicates that there is no sequence homology of this gene in *E. coli* with that in *Klebsiella pneumoniae* because *espB* is conserved at high frequency in *E. coli* as mentioned by (19). On the other hand *cesT/eae* is present in only 3 isolates as shown in fig (1).



Fig(1):-Gel electrophoresis of PCR of *cesT/eae* (chaperon type III secretion for *tir*/ intimate attachment to epithelial cell) amplicon product. L: ladder; 2, 3, 4,5,6,7,8: no. of isolates

There are several possible scenarios regarding the origin and evolution of the LEE. On the other hand, the observation that the organization and functions of the LEE are conserved among studied *E. coli* strains (20) and that the locus has a different G-C proportion to the rest of the chromosome (1) suggests that this group of genes was transferred as a cassette from some other bacterial species and was inserted as a whole in the *E. coli* chromosome. Also, there is increasing evidence of horizontal transfer of some genes of the LEE, recombination within this locus (21) as well as recombination with other pathogenic genes (22). These later observations suggest that this locus was assembled by multiple independent recombination events (23). the regulation of the *tir-cesT-eae* operon (henceforth referred as the *LEE5* operon), whose expression is directed by a promoter located upstream of the *tir* gene.


Also *cnf1* was detected amplification by using *cnf1* primer in *klebsiella* genome, it was observed that *cnf1* was exist in only 4 isolates as shown in fig (2). The presences of this gene may be attributed to horizontal gene transfer in which the genetic markers are transferred through many mechanisms mostly by conjugation or transformation .

So, according to that the production of *cnf1* will render such bacteria to grow and survive at the site of infection due to its ability to prevent the healing of wound through its effect on polymorphonuclear leukocytes.

(24) observed that *cnf1* is mostly found among enteric bacteria associated with urinary and digestive tract infections and the present study suggests that the source of *Klebsiella pneumoniae* isolates from wound infections due to contamination the wounds with urine or stool containing this bacteria.

The mechanisms of action of cytotoxic necrotizing factor-1(CNF-1) involve the Rho-dependent rearrangement of the cytoskeleton in eukaryotic cells with a complex of consequences (12).

L 2 3 4 5 6 7 8



Fig(2):-Gel electrophoresis of PCR of cytotoxic necrotizing factor-1 *cnf1* amplicon product. L: ladder; 2, 3, 4,5,6,7,8: no. of isolates

References:-

- 1- Elliott S J, Wainwright L A, McDaniel T K, Jarvis K G, Deng Y K, Lai L C, McNamara B P, Donnenberg M S, Kaper J B. The complete sequence of the locus of enterocyte effacement (LEE) from enteropathogenic *Escherichia coli* E2348/69. *Mol Microbiol.* 1998;**28**:1–4
- 2- Jarvis K G, Giron J A, Jerse A E, McDaniel T K, Donnenberg M S, Kaper J B. Enteropathogenic *Escherichia coli* contains a putative type III secretion system necessary for the export of proteins involved in attaching and effacing lesion formation. *Proc Natl Acad Sci USA.* 1995;**92**:7996–8000.
- 3- Abe A, de Grado M, Pfuetzner R A, Sánchez-SanMartín C, DeVinney R, Puente J L, Strynadka N C J, Finlay B B. Enteropathogenic *Escherichia coli* translocated intimin receptor, Tir, requires a specific chaperone for stable secretion. *Mol Microbiol.* 1999;**33**:1162–1174.
- 4- Friedberg D, Umanski T, Fang Y, Rosenshine I. Hierarchy in the expression of the locus of enterocyte effacement genes of enteropathogenic *Escherichia coli*. *Mol Microbiol.* 1999;**34**:941–952.
- 5- Elliot, S. J., Wainwright, L. A., McDaniel, T. K., Jarvis, K. G., Deng, Y. K., Lai, L. C., MacNamara, B. P., Donnenberg, M. S. & Kaper, J. B. (1998). The complete sequence of the locus of enterocyte effacement (LEE) from enteropathogenic *Escherichia coli* E2348/69. *Mol Microbiol* 28, 1-4.
- 6- Donnenberg, M. S., Yu, J. & Kaper, J. B. (1993). A second chromosomal gene necessary for intimate attachment of enteropathogenic *Escherichia coli* to epithelial cells. *J Bacteriol* 175, 4670-4680
- 7- Ebel, F., Deibel, C., Kresse, A. U., Guzman, C. A. & Chakraborty, T. (1996). Temperature- and medium-dependent secretion of proteins by shiga toxin-producing *Escherichia coli*. *Infect Immun.* 64, 4472-4479.
- 8- Abe, A., Kenny, B., Stein, M. & Finlay, B. B. (1997). Characterization of two virulence proteins secreted by rabbit enteropathogenic *Escherichia coli*, EspA and EspB, whose maximal expression is sensitive to host body temperature. *Infect Immun.* 65, 3547-3555.



- 9- Frankel, G., Philips, A. D., Rosenshine, I., Dougan, G., Kaper, J. B. & Knutton, S. (1998). Enteropathogenic and enterohaemorrhagic *Escherichia coli* : more subversive elements. *Mol Microbiol* 30, 911-921.
- 10- Foubister, V., Rossenshine, I., Donnenberg, M. S. & Finlay, B. B. (1994). The *eaeB* of enteropathogenic *Escherichia coli* is necessary for signal transduction in epithelial cells. *Infect Immun* 62, 3038-3040.
- 11- Kenny, B. & Finlay, B. B. (1997). Intimin dependent binding of enteropathogenic *Escherichia coli* to host cells triggers novel signaling events, including tyrosine phosphorylation of phospholipase C-1. *Infect Immun* 65, 2528-2536.
- 12- Doye, A., A. Mettouchi, G. Bossis, R. Clement, C. Buisson-Touati, G. Flatau, L. Gagnoux, M. Piechazyk, P. Boquet, and E. Lemich. (2002). CNF1 exploits the ubiquitin-proteasome machinery to restrict Rho GTPase activation for bacterial host cell invasion. *Cell*. 111:553-564.
- 13- Boquete, P. (1998). Cytotoxic necrotizing factor 1 *Escherichia coli*: a toxin with a new intracellular activity for eukaryotic cells. *Folia Microbiol.* 43:285-289.
- 14- Caprioli, A., V. Falbo, L.G. Roda, F. M. Ruggeri, and C. Zona. (1983). Partial purification and characterization of an *Escherichia coli* toxic factor that induces morphological cell alteration. *Infect. Immune*. 39: 1300-1306.
- 15- Bond, M., R. P. Fabunmi, A. H. Baker, and A. C. Newby. (1998). Synergistic upregulation of metalloproteinase-9 by growth factors and inflammatory cytokines: an absolute requirement for transcription factor NF-kappa B. *FEBS Lett.* 435: 29-34.
- 16- Mackay, D. J. G., Esch, F., Furthmayr, H., and Hall, A. (1997). Rho- and Rac-dependent assembly of focal adhesion complexes and actin filaments in permeabilized fibroblasts: an essential role for ezrin/radixin/moesin proteins. *J. Cell Biol.* 138,927-938.
- 17- Holt JC, Krieg NR, Sneath A, Stachley JT, William ST (1994). *Bergey's manual of determinative bacteriology*, 9th ed. USA, pp.552.
- 18- Yamamoto, S., Terai, A., Yuri, K., Kurazono, H., Takeda, Y. and O. Yoshida. (1995). Detection of urovirulence factor in *Escherichia coli* by multiplex polymerase chain reaction. *FEMS Immunol Med Microbiol.* 12:85-90
- 19- Frankel, G., Philips, A. D., Rosenshine, I., Dougan, G., Kaper, J. B. & Knutton, S. (1998). Enteropathogenic and enterohaemorrhagic *Escherichia coli* : more subversive elements. *Mol Microbiol* 30:911-921.
- 20- Beltrametti, F., Kresse, A. U. & Guzman, C. A. (1999). Transcriptional regulation of the *esp* genes of enterohaemorrhagic *Escherichia coli*. *J Bacteriol* 181: 3409-3418.
- 21- McGraw, E. A., Li, J., Selander, R. K. & Whittam, T. S. (1999). Molecular evolution and mosaic structure of *a*, *b*, and *c* intimins of pathogenic *Escherichia coli*. *Mol Biol Evol* 16: 12-22.
- 22- Lederberg, J. (1998). Emerging infection : an evolutionary perspective. *Emerg Infect Dis* 4, 366-371.
- 23- Reid, S. D., Herbelin, C. J., Bumbaugh, A. C., Selander, R. K. & Whittam, T. S. (2000). Parallel evolution of virulence in pathogenic *Escherichia coli*. *Nature* 406, 64-67.
- 24- Sare, T., Alessia, F. and Carla, F. (2008). The Rho activating CNF-1 toxin from pathogenic *Escherichia coli*: A risk factor for human cancer development. *J. Bio. Med.* 3:4.



الخلاصة:-

في هذه الدراسة، تم عزل وتشخيص سبع *Klebsiella pneumoniae* حيث تم التحري على احتواء عزلات من بكتريا العزلات للجينات *espB*, *cesT/eae*, and *cnf1*. وقد وجد ان جميع العزلات غير حاوية على *espB* في حين وجد ان ثلاث عزلات فقط حاوية على *cesT/eae*. كما وجد ان اربع عزلات فقط حاوية على *cnf1* جين. كما لوحظ ان وجود هذه الجينات في البكتريا يزيد من قابليتها على غزو وتجنب الجهاز المناعي ويزيد من قابليتها على حدوث الاصابة وبالأخص وجود جينات *cnf1*.