

# On Harmonic Index of Some Special Graphs with Certain Vertex Gluing Graphs

Haneen Kareem Aljanabi  
Faculty of Computer Science and Mathematics  
University of Kufa  
Al-Najaf, Iraq  
[Haneenk.aljanabi@uokufa.edu.iq](mailto:Haneenk.aljanabi@uokufa.edu.iq)

Received Dec. 18, 2017. Accepted for publication Jan. 16, 2018

DOI: <http://dx.doi.org/10.31642/JoKMC/2018/050202>

**Abstract:** The stability of cycloalkanes strain energy, branched alkanes and linear alkanes, can be reasonably modeled using the harmonic index ( $H$ ), which is commonly defined as  $H(G) = \sum_{u-v} \frac{2}{d(u)+d(v)}$  where  $d(u)$  denotes the vertex degree of  $u$  in graph  $G$ . In the current work, public equation is derived to the  $H$  index of vertex gluing and certain graphs.

**Keywords:** Harmonic index, Vertex Gluing.

## 1. INTRODUCTION


Characteristics of molecules can be physically/chemically modeled using tools called topological indices, which have the potentials to design the active pharmacological compounds, recognize the environmentally dangerous materials, etc. [2]. Therefore, authors have been attracted to present a variety of researches on such indices, to name a few. The harmonic index, which is considered of great importance, is utilized in modeling the branching of alkanes skeleton that includes carbon-atom. In the 1980s, a computer code to model the automatic generation of conjectures was established by Siemion Fajtlowicz using graph theory, where the likely relations between countless graph invariants, including the vertex-degree-based, were investigated [10-11]. Subsequently, however, no considerable attention has been paid for the  $H(G)$  predicted, especially by the chemists. So far, this quantity has not been reintroduced until 2012, when Zhang introduced it as a “harmonic index”, which has recently attracted many researchers to readopt this methodology. For example, the harmonic index was successfully employed in chemical applications, but, knowing the present situation in mathematical

chemistry, such authors are very much to be expected [1], [3-6], [9].


In this theory, the graph is symbolized as  $G=(V,E)$ , where  $V = \{v_1, \dots, v_{n-1}, v_n\}$  is the vertex set, while  $E=E(G)$  is the edge set of  $G$ . Assuming that  $v_i v_j \in E$ , hence,  $G - v_i v_j$  can be denote to left graph  $G$  after cancelling the edge  $v_i v_j$ . In a molecular graph, the valence of a molecule, i.e. bonds number on a vertex, is represented by the degree of each vertex. In case that  $d(v)$  equals to unity,  $v$  is said to be a pendent vertex. Moreover, the edge being incident to a pendent vertex is known as a pendent edge. Finally, a chemical is that sort of trees with maximal vertex degree does not exceed 4 [7].

## 2. BASIC DEFINITIONS AND KNOWN RESULTS


“A  $K_4$ –homeomorphism graph is the graph obtained when the six edges of a complete graph with four vertices ( $K_4$ ) are subdivided into  $(e_1, e_2, e_3, e_4, e_5, e_6)$  segments, respectively [8]” see “Fig. 1.”.




"A complete bipartite graph is a simple bipartite graph with partite sets  $V_1$  and  $V_2$ , where every vertex in  $V_1$  is adjacent with all vertices in  $V_2$ . If  $|V_1|=n$  and  $|V_2|=m$ , then such complete bipartite graph is denoted by  $K_{n,m}$  or  $(K(n,m))$ . So  $K_{n,m}$  has order  $(n+m)$  and size  $(nm)$ . A tree  $K_{1,m}$  is also called a star [8]" see "Fig. 2,".



"A  $(r)$  – bridge graph is a graph consisting of  $s$  paths joining two vertices, which is denoted by  $Q(a_1, \dots, a_r)$ , where  $a_1, \dots, a_r$ , are the lengths of  $r$  paths [8]" see "Fig. 3,".



"A web graph  $Web(n,m)$  is the graph obtained from the Cartesian product of the cycle  $C_n$  and the path  $P_m$  [8]" see "Fig. 4,".



Firstly, examples of some simple graphs are solved for the harmonic index.

*Lemma 2.1.*

Let  $K_n$  is a complete graph of order  $n$ , then the harmonic index,  $H(K_n) = \frac{n}{2}$ .

*Lemma 2.2.*

Let  $C_n$  be a cycle of order  $n$ ,  $n \geq 3$  and size  $m$  ( $n=m$ ), then the harmonic index of  $C_n$  is,  $H(C_n) = \frac{n}{2}$ .

*Lemma 2.3.*

Let  $S_n$  be a star of order  $n$ , then the harmonic index of  $S_n$ ,  $H(S_n) = \frac{2(n-1)}{n}$ .

*Lemma 2.4.*

Let  $P_n$  be a path of order  $n$ , then the harmonic index of a path  $P_n$  is as follows:

- 1- When  $n=2$  then  $H(P_1) = 1$ .
- 2- When  $n=3$  then  $H(P_3) = \frac{4}{3}$ .
3. When  $n \geq 4$  then  $H(P_n) = \frac{4}{3} + \frac{1}{2}(n-3)$

### 3. THE HARMONIC INDEX OF SOME SPECIAL GRAPHS

*Theorem 3.1.*

Let  $e_1, e_2, e_3, e_4, e_5$  and  $e_6$  be positive integers. The harmonic index of  $k_4$ -homeomorphism graph is denoted by  $K_4(e_1, e_2, e_3, e_4, e_5, e_6)$  as follows:

- 1- If  $e_1, e_2, e_3, e_4, e_5$  and  $e_6 = 1$ , then the H-index of  $K_4$  – homeomorphism is 2.
- 2- If  $e_1, e_2, e_3, e_4, e_5$  and  $e_6 \neq 1$  such that  $e_i, (i=1, \dots, 6)$  have  $k$  vertices in every  $e_i$ , then the H-index of  $K_4$  – homeomorphism is  $\frac{24}{5} + 3(k-1)$

*Proof:*

- 1- If  $e_1, e_2, e_3, e_4, e_5$  and  $e_6 = 1$ , thus each of them has two vertices of the same third degree besides only one edge. Thus, the H – index will be:

$H(K_4\text{-homeomorphism})$

$$= \sum_{u \sim v} \frac{2}{d(u) + d(v)} = 6 \cdot \frac{2}{3+3} = 2$$

2- If  $e_1, e_2, e_3, e_4, e_5$  and  $e_6 \neq 1$  see “Fig. 1,” then each one of them has two or more edges having three degrees for their vertices located at the terminals and only two degrees for the remaining vertices. Therefore, every  $e_i$  ( $i=1, \dots, 6$ ) will have  $(k-1)$  edge; and hence, all remaining  $(k-3)$  edges will be of the same two degree. So, we have  $6(k-3) \frac{2}{2+2}$  edges.

That is,

$$H(K_4(e_1, e_2, e_3, e_4, e_5, e_6)) = 12 \cdot \frac{2}{2+3} + \frac{12}{4}(k-3) = \frac{24}{5} + 3(k-3)$$

**Theorem 3.2.**

If  $n, m$  are positive integers, then the harmonic index of a web graph is denoted by  $\text{web}(n, m)$  as follows:

- 1- If  $m=2$  then,  $H(\text{web}(n, 2)) = n$ .
- 2- If  $m=3$  then,  $H(\text{web}(n, 3)) = \frac{125n}{84}$ .
- 3- If  $m \geq 4$ , then  $H(\text{web}(n, m)) = \frac{n[42m-1]}{84}$ .

*Proof:*

- 1- When  $m=2$  and  $n \geq 3$ , then the web  $(n, 2)$  is obtained from the Cartesian product of the  $C_n$  and  $P_2$  see “Fig. 5 (a).” Thus, we have  $2n$  edges, where each one of them have two vertices. Then,  $d(v_i)=3$ ; ( $i=1, 2$ ). Since every path in the web graph is incident with the vertex of  $C_n$  of  $n \geq 3$ , then we have  $n$  paths with one edge for each of them, So, we have  $n$  edges with two vertices for each one of them, where every vertex  $d(u_i) = 3$ ; ( $i=1, 2$ ). Hence, by the definition of harmonic index:

$$H(\text{web}(n, 2)) = 2n \frac{2}{3+3} + n \frac{2}{3+3} = \frac{4n+2n}{6} = n$$


- 2- When  $m=3$  and  $n \geq 3$  then web  $(n, 3)$  is obtained from  $(C_n)$  and  $(P_3)$  see “Fig. 5.(b).” So, we have  $2n$  edges obtained from the edges of first cycle and third cycle, where each one of them have two vertices. Thus,  $d(v_i)=3$ ; ( $i=1, 2$ ). If we have

$(n)$  edges (the number of edges for the second cycle), where each one of them have two vertices  $d(v_i) = 4$ ; ( $i=1, 2$ ). Also, we have  $n$  paths, each path have 2 edges with two vertices for each edge, then,  $d(u_1) = 3$  and  $d(u_2) = 4$ . Hence, by definition of harmonic index:

$$H(\text{web}(n, 2)) = 2n \frac{2}{3+3} + n \frac{2}{4+4} + 2n \frac{2}{3+4} = \frac{125n}{84}$$

- 3- When  $m \geq 4$  and  $n \geq 3$ , then the web  $(n, m)$  is the web graph calculated from  $C_n$  and  $P_m$  see “Fig. 5 (c).” Thus, we have  $(2n)$  edges obtained from the edge of first cycle and final cycle, where each one of them have two vertices as  $d(v_i) = 3$ ; ( $i=1, 2$ ). Then,  $(m-2)n$  edges of the total number of cycle edges “excluding the first and final cycles” will have two vertices with  $d(v_i) = 4$ ; ( $i=1, 2$ ). Also, we have  $n$  paths with  $(m-1)$  edges for each of them, where two edges in every path are incident with two vertices of degree 3 and 4. Thus,  $d(u_1)=3$  and  $d(u_2)=4$ . Moreover,  $m-3$  edges in every path are incident with two vertices of degree 3; that is  $d(u_i)=3$  and ( $i=1, 2$ ). subsequently, the harmonic index definition is,

$$\begin{aligned} H(\text{web}(n, m)) &= 2n \frac{2}{3+3} + (m-2)n \frac{2}{4+4} + 2n \frac{2}{3+4} \\ &+ (m-3)n \frac{2}{4+4} \\ H(\text{web}(n, m)) &= \frac{2n}{3} + \frac{2n(m-2)}{8} + \frac{4n}{7} + \frac{n(m-3)}{4} \\ &= \frac{2n}{3} + \frac{nm}{8} - \frac{n}{2} + \frac{4n}{7} + \frac{nm}{4} - \frac{3n}{4} \\ &= \frac{56n - 42n - 63n + 18}{84} + \frac{1}{2}nm \\ &= \frac{-n}{84} + \frac{nm}{2} = \frac{42nm - n}{84} = \frac{n[42m-1]}{84} \end{aligned}$$




Suppose that  $n, m$  are positive integers, then the complete bipartite graph, which is denoted by  $(K_{n,m})$ , will have a harmonic index of  $H(K_{n,m}) = \frac{2nm}{n+m}$

*Proof:*

If we have a number of edges  $nm$ , where every one of these edges has couple of vertices having equal degree see “Fig. 6,” i.e.  $m$  and  $n$  are respectively the degree of first and second vertex. Hence, by definition of harmonic index, we get

$$H(K_{n,m}) = \sum_{u-v} \frac{2}{d(u)+d(v)} = nm \frac{2}{n+m} = \frac{2nm}{n+m}$$



“Fig. 6. complete bipartite graph”

**Theorem 3.4.**

Let  $Q(a_1, a_2, \dots, a_k)$  be a  $k$  – Bridge graph, ( $k$ ) is a positive integer, and  $n$  is the number of vertices for any path. Then, the harmonic index of the  $k$  – Bridge is:

$$H(Q(a_1, a_2, \dots, a_k)) = \frac{4k}{k+2} + \frac{k}{2}(n-3)$$


*Proof:*

Let  $k=3$ , then the graph  $G=Q(a_1, a_2, a_3)$  is as shown in “Fig. 7,” : Thus,

$$H(Q(a_1, a_2, a_3)) = \frac{12}{5} + \frac{3}{2}(n-3)$$

Hence, it is true with,

$$\begin{aligned} H(Q(a_1, a_2, \dots, a_k)) &= \frac{4k}{k+2} + \frac{k}{2}(n-3) \\ &= \frac{4 \cdot 3}{3+2} + \frac{3}{2}(n-3) = \frac{12}{5} + \frac{3}{2}(n-3); \text{ when } k=3. \end{aligned}$$




“Fig. 7. (3) – Bridge graph  $Q(a_1, a_2, a_3)$ ”

Assuming that this theory is valid for  $k=r$  ( $r \geq 3$ ), then the H-index for  $Q(a_1, a_2, \dots, a_r)$  is given by:

$$H(Q(a_1, a_2, \dots, a_r)) = \frac{4r}{(r+2)} + \frac{r}{2}(n-3).$$

Constructing the graph  $(Q_{r+1})$  as follows,  $Q(a_1, a_2, \dots, a_r)$  has the sample:

Where the location of the edges represents  $a_1$  included within graph  $Q(a_1, a_2, \dots, a_r)$  at the  $i^{\text{th}}$  position see “Fig. 8,”.




“Fig. 8. (r) – Bridge graph”

The graph  $U$  is the path that contains terminals  $u_1$  and  $u_n$ , where  $n$  is the number of vertices in  $U$  as follows see “Fig. 9,”.



“Fig. 9. Path”

Connecting the graph  $Q(a_1, a_2, \dots, a_r)$  with the graph ( $U$ ) such that  $v_1 = u_1$  and  $v_n = u_n$ , then, the vertices  $v_1 = u_1$  and  $v_n = u_n$  are of degree  $(r+1)$  see “Fig. 10,”.



“Fig. 10. (r+1) – Bridge graph”

Thus,

$$H(Q_{r+1}) = \frac{4r}{(r+1)+2} + \frac{r}{2}(n-3) + \frac{4}{(r+1)+2} + (n-3) \cdot \frac{1}{2}$$


$$H(a_1, a_2, \dots, a_{r+1}) = \frac{4}{(r+3)}(r+1) + \frac{n-3}{2}(r+1),$$

Since  $(k=r+1)$

$$\begin{aligned} H(a_1, a_2, \dots, a_{r+1}) &= \frac{4k}{k+2} + \frac{k}{2}(n-3) \\ &= \frac{4k}{k+2} + \frac{k}{2}(n-3) \end{aligned}$$

#### 4. THE HARMONIC INDEX OF CERTAIN VERTEX GLUING GRAPHS

“Let  $v_1$ -gluing of Complete bipartite graph be a graph obtained from two different Complete bipartite graphs  $K_{n,m}$  and  $K_{h,t}$  with common one vertex  $v_1$  denoted by  $K_{n,m}^{h,t}(v_1)$  (vertex gluing of graph) [8]”, see “Fig. 11,”



“Fig. 11. ( $v_1$ ) – gluing of Complete bipartite graph”

**Theorem 4.1.**

Suppose that  $n, m, h$  and  $t$  are positive integers, then, the  $v_1$ - gluing of a complete bipartite graph  $H$  –index  $K_{n,m}^{h,t}(v_1)$  is:

$$H(K_{n,m}^{h,t}(v_1)) = \frac{2m(n-1)}{(n+m)} + \frac{2t(h-1)}{(h+t)} + \frac{2m}{(n+m+t)} + \frac{2t}{(h+m+t)}$$

*Proof:*

We have two complete bipartite graphs connected with a common vertex, where every one of them has  $H$  – index  $\frac{2nm}{n+m}$  by using Theorem 3.3. Thus,


deleting the  $H$  – index of the  $m$  edges connected with  $v_1$ - gluing equals to  $(m \frac{2}{n+m})$ . On the other hand, adding the  $H$ -index of the  $m$  edges equals to  $(m \frac{2}{n+(m+t)})$ . Thus, every edge have degrees of  $n$  and  $(m+t)$ , when  $(m+t)$  is the degree of vertex  $v_1$  – gluing. So,

$$H(K_{n,m}^{h,t}(v_1)) = \frac{2nm}{(n+m)} - m \frac{2}{(n+m)} + m \frac{2}{n+(m+t)} + \frac{2ht}{(h+t)} - t \frac{2}{(h+t)} + t \frac{2}{h+(m+t)}$$

$$= (nm-m) \frac{2}{(n+m)} + (ht-t) \frac{2}{(h+t)} + \frac{2m}{n+(m+t)} + \frac{2t}{h+(m+t)}$$

$$= \frac{2m(n-1)}{(n+m)} + \frac{2t(h-1)}{h+t} + \frac{2m}{n+m+t} + \frac{2t}{h+m+t} \quad \blacksquare$$

“Let  $K_4^2$  – homeomorphism be a graph obtained from two deferent  $K_4$  – homeomorphism graphs  $K_4(e_1, e_2, e_3, e_4, e_5, e_6)$  and  $K_4(e_7, e_8, e_9, e_{10}, e_{11}, e_{12})$ , with common one vertex  $v_1$  (vertex gluing of graph) [8]”, see “Fig. 12,”



“Fig. 12.  $K_4^2$  – homeomorphism graph”

**Theorem 4.2.**

Let  $e_i$  be a positive integer such that  $1 \leq i \leq 12$  and every  $e_i$  has  $k$  vertices and an edge length of  $(k-1)$ , then the harmonic index of the  $K_4^2$  – homeomorphism graph is:

1- If length of  $e_i=1$  then,


$$H(K_4^2 \text{ – homeomorphism}) = \frac{10}{3}.$$

2- If length of  $e_i=2$  see “Fig. 13,” then,

$$H(K_4^2 \text{ – homeomorphism}) = \frac{87}{10}.$$

3- If length of  $e_i=k-1, k \geq 4$  then,

$$H(K_4^2 \text{ – homeomorphism}) = \frac{87}{10} + 6(k-3).$$



“Fig. 13.  $K_4^2$  – homeomorphism graph ( $e_i=2$ )”

*Proof:*

1- By using the definition,  $\sum \frac{2}{d(u)+d(v)} = \frac{10}{3}$

2- By using the definition then,

$$H(K_4^2 \text{ – homeomorphism}) = \frac{87}{10}$$

3- We prove that by mathematical indication:

If  $k=4$ , then the length of  $e_i=3$ . Using the definition, we

$$\text{get } H(K_4^2 - \text{homeomorphism}) = \frac{147}{10}$$

It is true that,

$$H(K_4^2 - \text{homeomorphism}) = \frac{87}{10} + 6(k-3)$$

$$= \frac{87}{10} + 6(4-3) = \frac{147}{10}$$

Assuming that this is true for  $r \geq 4$  see "Fig. 12," so,

$$H(K_4^2 - \text{homeomorphism}) = \frac{87}{10} + 6(r-3).$$


To prove this is true for  $k=r+1$ , a one vertex is added for all, i.e.  $e_i$ ,  $i=1, 2, \dots, 12$ . So, the number of edges is increased (1) in every  $e_i$ , that is we have a new edge  $a_i$  in every  $e_i$  see "Fig. 14,". Two vertices for  $a_i$  have 2 degrees, and we have 12 edges for  $K_{42}$ -homeomorphism.

Hence, we add  $12 \cdot \frac{2}{2+2}$  for  $\frac{87}{10} + 6(r-3)$ , that is,

$$H(K_4^2 - \text{homeomorphism}) = \frac{87}{10} + 6(r-3) + 12 \cdot \frac{2}{2+2}$$


$$= \frac{87}{10} + 6(k-3) + 6$$

$$= \frac{87}{10} + 6[r-3+1] \text{ since } k=r+1, = \frac{87}{10} + 6[k-3]$$



"Fig. 14:  $K_4^2$ -homeomorphism graph ( $e_i=1,2,\dots,r+1$ )"

"Let  $v_1$ -gluing of  $k, t$  - Bridge graph be a graph obtained from two different  $k$  - Bridge graph  $Q_1$  and  $t$ -Bridge graph  $Q_2$  with common vertex  $v_1$  denoted by  $Q_k^t(v_1)$  (vertex gluing of graph) [8]", see "Fig. 15,".



"Fig. 15. ( $v_1$ ) - gluing of ( $k, t$ ) - Bridge graph"

**Theorem 4.3.**

Let  $k, t, s$  and  $h$  be positive integers. Then, the  $v_1$  - gluing of  $k, t$  - Bridge graph of  $H$  - index  $Q_k^t(v_1)$  is:

$$H(Q_k^t(v_1)) = \frac{2k}{k+2} + \frac{2t}{t+2} + \frac{2(k+t)}{k+t+2} + \frac{k(n-3)}{2} + \frac{t(h-3)}{2}$$

**Proof:**

We have two  $k$ - Bridge graph connected with common vertex  $v_1$  see "Fig. 15," and every vertex of them have  $H$ -index  $\frac{4k}{k+2} + \frac{k(n-3)}{2}$ . By Theorem 3.4,

deleting the  $H$ -index for  $k$  edges connected with  $v_1$ - gluing deleting  $k \cdot \frac{2}{k+2}$ , and adding the  $H$ -index of  $k$

edges  $k \cdot \frac{2}{2+(k+t)}$  that is every edge have degree 2 and

$(k+t)$ , where  $(k+t)$  is the degree of the vertex  $v_1$  - gluing. Similarly to what followed with  $t$  - Bridge,


$$H(Q_k^t(v_1)) = \frac{4k}{k+2} + \frac{k(n-3)}{2} + \frac{4t}{t+2} + \frac{t(h-3)}{2}$$

$$- \frac{2k}{t+2} - \frac{2t}{t+2} + \frac{2k}{(k+t)+2} + \frac{2t}{(k+t)+2}$$

$$= \frac{2k}{k+2} + \frac{2k}{k+t+2} + \frac{k(n-3)}{2} + \frac{2t}{t+2} + \frac{2t}{k+t+2} + \frac{t(h-3)}{2}$$

$$= \frac{2k}{k+2} + \frac{2t}{t+2} + \frac{2(k+t)}{k+t+2} + \frac{k(n-3)}{2} + \frac{t(h-3)}{2}$$

"Let  $v_1$  - gluing of web graph be a graph obtained from two different web graphs web  $(n, m)$  and web  $(s, t)$  with one common vertex  $v_1$  denoted by  $W_{n,m}^{s,t}(v_1)$  (vertex gluing of graph) [8]" see "Fig. 16,".



"Fig. 16.  $v_1$ - gluing of web graph"

**Theorem 4.4.**

Let  $(n, m, s$  and  $t)$  be positive integers. Then, the  $v_1$ - gluing of web graph of  $H$ -index  $W_{n,m}^{s,t}(v_1)$  is:

1. If  $m, s=2$  and  $n, t \geq 3$  then,

$$H(W_{n,2}^{t,2}(v_1)) = n + t - \frac{2}{3}$$

2. If  $m, s=3$  and  $n, t \geq 3$  then,

$$H(W_{n,3}^{t,3}(v_1)) = \frac{125}{84}(n+t) - \frac{194}{315}$$

3. If  $m, s \geq 4$  and  $n, t \geq 3$  then,

$$H(W_{n,m}^{t,s}(v_1)) = \frac{3n[42m-1] + 3t[42s-1] - 168}{84}$$

*Proof:*

We have two web graphs connected with a common vertex  $v_1$ , where every vertex of them has an H-index as:

- 1-  $H(\text{web}(n,2)) = n$  by Theorem 3.2 (1), we delete

$3 \cdot \frac{2}{3+3}$ , which are the edges connected with  $v_1$ , and

we add  $3 \cdot \frac{2}{3+6}$ , which are the edges connected with

$v_1$ -gluing see "Fig. 17 (a)". Similarly, work to  $H(\text{web}(t,2))$  so,

$$\begin{aligned} H(W_{n,2}^{t,2}(v_1)) &= n + t - \frac{2}{3} = n - 3 \cdot \frac{2}{3+3} + 3 \cdot \frac{3}{3+6} \\ &+ t - 3 \cdot \frac{2}{3+3} + 3 \cdot \frac{3}{3+6} = n + t - \frac{2}{3} \end{aligned}$$

This proof is similar to proof (1), but one of the deleted edges has 3 degrees on one vertex and 4 on the other vertex, while one of the added edges has (4) degrees on one vertex and 6 on the other vertex see "Fig. 17 (b)". We use Theorem 3.2 (2).

$$\begin{aligned} H(W_{n,3}^{t,3}(v_1)) &= n + t - \frac{2}{3} = \frac{125n}{84} + \frac{125t}{84} - 2 \cdot \frac{2}{3+4} \\ &- 4 \cdot \frac{2}{3+3} + 2 \cdot \frac{2}{6+4} + 4 \cdot \frac{2}{6+3} = \frac{125(n+t)}{84} - \frac{194}{315} \end{aligned}$$





Fig. 17.  $v_1$ -gluing of web graph  $\text{web}(t,2)$ , and  $\text{web}(t,3)$

- 3- This proof is similar to proof (2) and by Theorem 3.2 (3) then,

$$\begin{aligned} H(W_{n,m}^{t,s}(v_1)) &= \frac{n[42m-1]}{84} + \frac{t[42s-1]}{84} - \frac{194}{315} \\ &= \frac{(n+t)[42m-1+42s-1]}{84} - \frac{194}{315} \\ &= \frac{(n+t)[42(m+s)-2]}{84} - \frac{194}{315} \end{aligned}$$

## References

- [1] A. Behtoei, M. Jannesari, B. Taeri, "Maximum Zagreb index, minimum hyper-Wiener index and graph connectivity," Appl. Math. Lett. 22 (2009) 15711576. <http://dx.doi.org/10.1016/j.aml.2009.05.001>.
- [2] K. C. Das, I. Gutman, B. Furtula, "Survey on Geometric Arithmetic indices of graphs," MATCH Commun. Math. Comput. Chem. 65 (2011) 595644.
- [3] M. V. Diudea, I. Gutman, L. Jntschi, "Molecular Topology," Nova, Huntington, 2002.
- [4] B. Liu, I. Gutman, "On general Randic indices," MATCH Commun. Math. Comput. Chem. 58 (2007) 147154.
- [5] B. Liu, I. Gutman, "On a conjecture on Randic indices," MATCH Commun. Math. Comput. Chem. 62 (2009) 143154.
- [6] B. Liu, Z. You, "A survey on comparing Zagreb indices," MATCH Commun. Math. Comput. Chem. 65 (2011) 581593.
- [7] M. A. Mohammed, K.A. Atan, A.M. Khalaf, M. R, Said, R. Hasni, "Atom Bond Connectivity Index Of Molecular Graphs of Alkynes and Cycloalkynes," Journal Of Combinatorial and Theoretical Nanoscience, Vol. 13, pp. 157-170, 2016.s

- [8] M. A. Mohammed, K.A. Atan, A.M. Khalaf, M. Rushdan, R. Hasni5, "The atom bond onnectivity index of certain graphs," International Journal of Pure and Applied Mathematics Volume 106 No. 2 2016, 415-427.
- [9] B. Zhou, N. Trinajstic, "Further results on atombond connectivity index of trees," Chem. Phys. Lett. 455 (2008) 120123.  
<http://dx.doi.org/10.1016/j.cplett.2008.02.064>.
- [10] L. Zhong, "The harmonic index on unicyclic graphs," Ars Combinatoria, Vol. 104, pp. 261-269, 2012.
- [11] L. Zhong, "The harmonic index for graphs", Applied Mathematics Letters, Vol. 25, No.3, pp. 561-566, 2012.