

Evaluation Of Immune Response To Brucella Infection In Sheep

Kareem Mohamed* Abdullah O Alhatami** Akram Motlak***

*Department of Community Medicine, Faculty of Medicine, University of Kufa

**Department of Microbiology, Faculty of Veterinary Medicine, University of Kufa

*** Department of Public health, Faculty of Veterinary Medicine, University of Kufa

Corresponding: Abdullah O Alhatami,

e-mail: abdullaho.mansour@uokufa.edu.iq

Abstract

Brucellosis in sheep and goats is a zoonotic infection (excluding *Brucella ovis*) with important effects on both public health and animal health. The disease is widespread in many areas of the world, particularly in some Mediterranean and Middle Eastern countries.

Aim of the study: the present study aims to find out the prevalence of brucellosis in sheep in one of the areas of An Najaf El-Ashraf city, and study of the immune response by measuring the levels of and interleukin-10 (IL-10).

Methodology: Current study conducted in the Najaf province (Al-Barakya) during the period between September 2013- March 2014, fulfilled by using Rose Bengal and Enzyme Immunosorbent Assay (ELISA) tests for estimation of brucella antibodies. A total number of 48 serum samples were collected from sheep and stored at -20°C.

Results: the frequency of positive sera by Rose Bengal test (RBT) was 20% and by ELISA technique was 25%. The level of interleukin 10 (IL-10) was estimated in all surveyed animals, 12 sera samples were showed an elevated IL-10 levels, all were positive with ELISA serological test.

Conclusion: the detection of significant concentration of IL-10 among the seropositive cases may be indicated an active infection of brucella with type 2 cytokine production.

Recommendations: the evaluation of immune response of vaccinated sheep with brucella REV1 vaccine, depending on the estimation of IFN- γ and IL-10 ratio that will reflect the cytokines induced during vaccination and consequently, the effective or not immune response.

Key word:

Brucellosis, sheep, Rose Bengal, ELISA, Interleukin 10

تقييم الاستجابة المناعية لداء البروسيليا في الأغنام

* كريم غالي محمد ** عبدالله عبيس الحاتمي *** اكرم مطلق صويح

* فرع طب المجتمع، كلية الطب/جامعة الكوفة ** فرع الاحياء المجهرية، كلية الطب البيطري/ جامعة الكوفة
*** فرع الصحة العامة، كلية الطب البيطري/ جامعة الكوفة**الخلاصة:**

يعتبر داء البروسيلات في الأغنام والماعز من الامراض المشتركة (عدا البروسيليا الغنمية) وله تأثيرات هامة على الصحة العامة والصحة الحيوانية هو مرض واسع الانتشار ويعتبر من الامراض المعدية ذات الاهمية. المرض منتشر في مناطق كثيرة من العالم، لا سيما في بعض مناطق حوض البحر الأبيض المتوسط والشرق الأوسط.

اهداف الدراسة: تهدف هذه الدراسة إلى معرفة مدى انتشار الاجسام المناعية لمرض البروسيليا في الأغنام في واحدة من المناطق من مدينة النجف الأشرف، ودراسة الاستجابة المناعية عن طريق قياس مستويات انترلوكين 10 (IL-10).

المنهجية: اجريت الدراسه الحاليه في محافظه النجف الاشرف وتحديدًا في منطقه البراكيه للفتره ما بين ايلول 2013 و آذار 2014 باستخدام تقنيات الاليزا و كاشف (Rose Bengal test) لتقدير مستويات الاجسام المضاده للبروسيليا. فاخذت 48 نماذج من امصال اغنام بطريقتي معقمتين فصلت بالطرد المركزي وتم تخزينها تحت درجة حراره (-20) لحين اجراء الفحوصات.

النتائج: اظهرت النتائج ان النتائج الايجابيه لكاشف (Rose Bengal test) هو (20%) , و (25%) بالنسبه لاختبار المقاييس المناعية المرتبطة بالانزيم. اخيرا تم قياس مستوى (Interleukin 10) في تلك الامصال والتي اظهرت ارتفاع مستواه في 12 نموذج وجميعها مترافقه مع نتائج ايجابيه بالنسبه لاختبار المقاييس المناعية المرتبطة بالانزيم.

الاستنتاج: ان الكشف عن تراكيز مرتفعة من الانترليوكين 10 في امصال الحيوانات الموجبة لاختبارات الكشف عن الاجسام المضاده للبروسيليا بتقنية المقاييس المناعية المرتبطة بالانزيم يمكن ان تؤثر اصابة نشطة بالبروسيليا مع وجود سايتوكينات من النوع الثاني.

التوصيات: توصي هذه الدراسة بتقييم الاستجابة المناعية للتطعيم الأغنام بلقاح البروسيليا REV1، اعتمادا على تقدير IFN- γ ونسبة IL-10 التي من شأنها أن تعكس السيتوكينات التي يسببها أثناء التلقيح وبالتالي، معرفة اذا كامن الاستجابة المناعية فعالة أم لا.

الكلمات المفتاحية: داء البروسيليا، الأغنام، فحص الرز بنغال، الاليزا، انترليوكين 10

Introduction:

Brucellosis in sheep and goats is a zoonotic infection (excluding *Brucella ovis*) with important effects on both public health and animal health. The disease is widespread in many areas of the world, particularly in some Mediterranean and Middle Eastern countries (1, 2).

Brucellosis in small ruminants is caused mainly by *Brucella melitensis*, which was the first species in the genus *Brucella* described. It is the most virulent one and most widely encountered of all the species (3). *Brucella melitensis* infection may cause abortion in pregnant animals or orchitis and epididymitis in adult males of sheep, goat and cow which may result in infertility (4). The number of

aborted animals is progressively increased in Middle East countries including Iraq during the last decades (5).

In Iraq, the prevalence of brucellosis in sheep was 0.9% in 1979, while several seroprevalence studies conducted in the recent decades showed that the prevalence of brucellosis in sheep and goat was increased markedly, in 2005 the prevalence was 8.59% (6).

Brucella spp. are facultative intracellular pathogens which resist killing by neutrophils, replicate inside macrophages and in "non-professional" phagocytes and maintain a long lasting interaction with the host cells. Therefore, host control of infection

requires a set of cells and factors like CD4+ and CD8+ T lymphocytes, T-helper 1(Th1) type cytokines such as (IFN γ) and TNF α , and activated macrophages and dendritic cells (DC) which together promote a complex response against *Brucella*. It has been postulated that Th1 cytokines contribute to control brucella infection (7, 8). Interleukin-12, chiefly a product of antigen-presenting cells (APC), is usually critical for the development of Th1 responses (9, 10). IL-10 is an anti-inflammatory cytokine secreted by T cells and macrophages. It interacts with the IL-10 receptor and like IFN- γ , signals through the Jack/Stat signaling pathway (11). It is known to downregulate Th1 response during Bucellosis (12).

The present study aims to find out the prevalence of brucellosis in sheep in one of the areas of An Najaf El-Ashraf city, and study of the immune response by measuring the levels of and interleukin-10 (IL-10) by ELISA. IL-10 level can be used as a criterion to determine the ability of immune response in controlling the disease.

Material and Methods:

Forty eight blood samples were collected from unvaccinated three flocks of sheep in Al-Barakya, Najaf province; a venipuncture of jugular vein was done under sterile condition to collect 10 ml of peripheral blood sample. Samples were conveyed to the laboratory in Najaf Veterinary Hospital for centrifugation to obtain the sera. The tubes were centrifuged at 3000 r.p.m., for 5 minutes, then the sera were collected in a small vials (eppendorf tubes) and stored in freezer at -20 till the time of use. Serum specimens were analyzed in two phases, In the first phase, all specimens were screened by the Rose Bengal test

(RBT) for brucella antibodies, second phase all specimens were screened by indirect ELISA test to detect brucella lipopolysaccharide (LPS) antibodies, and estimation of level of IL-10 by ELISA.

Rose Bengal test (RBT):

RBT was carried out according to (13) with brucella abortus S99 antigen (Spinreact SA, Girona, Spain). Briefly, 30 of μ L antigen was mixed on a white glossy ceramic tile, with an equal volume of sheep serum. The tile was then rocked at room temperature for 4 minutes and any visible agglutination and/or the appearance of a typical rim was taken as a positive result.

Indirect ELISA:

The SERELISA®Brucella OCB Ab Mono indirect kit (SYNBIOTICS EUROPE SAS, Cedex, France) was used to test the serum samples for antibodies to *B. abortus* and *B. melitensis* according to the manufacturer's instructions. The optical density (OD) values for each of the controls provided in the kit and serum samples in the wells were read at 450 nm using a microplate photometer (Universal Microplate Reader, Bio-Tek Instruments, Inc.).

IL-10 detection by ELISA:

Serum levels of IL-10 was measured using ELISA Kit supplied by (USBiological, USA). The assay was done according to protocol provided by the manufacturer.

Results:

Out of 48 serum samples collected from sheep revealed 10 (20.86%) were positive with Rose Bengal test, while 12 (25%) samples were positive indirect ELISA test.

Plotting standard curve for IL-10:

Standard preparation is necessary for immunoassay. Using a standard preparation, we draw a standard curve from graded reaction results of various standard

concentrations, and by comparison of a sample reaction result with the standard curve, we get assay value of the sample as in Figure 1.




Figure1. Standard curve for various standard concentrations

As shown in table 1, 12 out of 48 sera samples showed detectable levels of IL-10, all were ELISA positive.

Table 1. Serume IL-10 levels in study groups.

Sample number	O.D.	Concentration (Pg/ml)
Standard	0.368	2
Standard	0.348	1
Standard	0.321	0.75
Standard	0.317	0.375
Standard	0.267	0
1	0.247	0
2	0.401	2.22801
3	0.328	0.89442
4	0.484	3.74429
5	0.443	2.99528
6	0.402	2.22803
7	0.480	3.70015
8	0.404	2.22901
9	0.301	0.88242
10	0.309	0.88541
11	0.492	3.82272
12	0.490	3.81163
13	0.320	0.8890
14	0.114	0
15	0.211	0
16	0.122	0
17	0.212	0
18	0.112	0
19	0.211	0

20	0.115	0
21	0.186	0
22	0.211	0
23	0.118	0
24	0.112	0
25	0.162	0
26	0.117	0
27	0.192	0
28	0.187	0
29	0.114	0
30	0.193	0
31	0.161	0
32	0.217	0
33	0.114	0
34	0.115	0
35	0.201	0
36	0.115	0
37	0.113	0
38	0.212	0
39	0.147	0
40	0.265	0
41	0.157	0
42	0.242	0
43	0.276	0
44	0.211	0
45	0.165	0
46	0.123	0
47	0.122	0
48	0.112	0

Table 2 revealed that concentration of IL10 is significantly higher among those with positive test for brucellosis than those with negative test for brucellosis.

Table 2. Concentration of IL10 among those with or without brucellosis

	Number	(%)	Mean Conc. Of IL-10 (Pg/ml)
Brucellosis (+ve)*	12	(25%)	2.359198
Brucellosis (-ve)	36	(75%)	0

P value ≤ 0.05

***Brucella (+ve) = ELISA positive cases.**

Discussion:

Brucella melitensis is primarily responsible for brucellosis in sheep and goats, and the most important zoonotic agent among *Brucella* species (14). Small ruminant brucellosis remains a problem in some of industrialized countries as well as in all developing countries. Essentially, brucellosis almost always present where small ruminants are kept (15).

Rose Bengal test is internationally recommended for the screening of brucellosis in small ruminants (16). The standardization condition of the antigen limit the sensitivity of the test resulting in reduced performance for the diagnosis of *Brucella melitensis* infection in sheep (17, 18).

Present study revealed that the prevalence rate of brucellosis among sheep was higher than a previous report conducted in Iraq by (19) and (20) in Jordan. However, many other researchers in the Middle East and Mediterranean countries (have reported the seroprevalence rate to be between 26.66%- 27.1% in sheep and 27.7% in goat flocks (21, 22, 23), the present results indicating that the brucellosis is an endemic health problem of sheep the increased seroprevalence of brucellosis in Najaf could be due to illegal movement and trade of flocks of sheep (small ruminant) that regarded as a reservoir for brucella infection. however, the small sample size and selection of one region in Najaf province could not be representing the whole epidemiological picture in the city. Al-Hankawe and Rhaymah (24) mentioned that the total prevalence of brucellosis in Ninewah province was 11.8% using Rose Bengal test but the values varied markedly according to the areas, the highest prevalence was in

Al-Shekan (22.7%) among seven districts of the province.

The difference between percentage of seropositivity by ELISA and Rose Bengal could be due to the high sensitivity and specificity of ELISA test (13, 2).

IL-10 is a cytokine that regulates the balance between pathogen clearance and immunopathology. Previous findings demonstrate that IL-10 modulates the proinflammatory immune response to *B. abortus* infection and the lack of IL-10 increases resistance to *Brucella* infection (25).

It was obvious in the present findings that there was an increase in IL-10 level in the sera of animals having ELISA positive results.

Conclusion:

The detection of significant concentration of IL-10 among the seropositive cases may be indicated an active infection of brucella with type 2 cytokine production and could be down-regulation of type 1 cytokine function.

Recommendations:

The present study recommends the evaluation of immune response of vaccinated sheep with brucella REV1 vaccine, depending on the estimation of IFN- γ and IL-10 ratio that will reflect the cytokines induced during vaccination and consequently, the effective or not immune response.

References:

- 1- Abo-Shadi MA, Al-Harbi AIH, Ballal EM (2010) Serum Levels of Interferon Gamma in Patients with Brucellosis in a Saudi Hospital. *British Microbiology Research Journal* 4(3): 293-305.
- 2- Al-Hankawe OKh (2009) Effect of *Br. melitensis* strain vaccine (Rev.1) in titers of antibodies and interference with serological tests in

- ewes. Iraqi Journal of Veterinary Science, 23 supplement(1)Arabic:149-154.
- 3- Motamedi H, Darabpour E, Gholipour M, Seyyed Nejad, SM (2010) In vitro assay for the anti-brucella activity of medicinal plants against tetracycline-resistant *Brucella melitensis*. Journal of Zhejiang University SCIENCE B, Vol11 (7): 506-511.
 - 4- Vitry MA, Hanot Mambres D, De Trez C, Akira S, Ryffel B, Letesson JJ, Muraille E (2014) Humoral immunity and CD4+ Th1 cells are both necessary for a fully protective immune response upon secondary infection with *Brucella melitensis*. J. Immunol. 192:3740–3752.
 - 5- Obi TU, Amin I, Shareef JM, Hawez N, Abdi A (2000). Brucellosis. Veterinary Emergency plan, Northern Governorates of Iraq : 4-16. FAO NorthCoordination, Erbil.
 - 6- Sharief DM, Saleem HM, Al-Kubaisi AH, Mahdi AJ, Mohmood TS, Saeed EA, Al-Adhad BN (2006) Survey of the seroprevalence of Brucellosis in ruminants in IRAQ. International Symposia on Veterinary Epidemiology and Economics proceedings, ISVEE 11: Proceedings of the 11th Symposium of the International Society for Veterinary Epidemiology and Economics, Cairns, Australia, Theme 2 - Disease distribution & determinants: Poster session session, p 848.
 - 7- Copin R, De Baetselier P, Carlier Y, Letesson JJ, Muraille E (2007) MyD88-dependent activation of B220-CD11b+LY-6C+ dendritic cells during *Brucella melitensis* infection. J. Immunol. 178: 5182–5191.
 - 8- Fernandes DM, Jiang X, Jung JH, Baldwin CL (1996) Comparison of T cell cytokines in resistant and susceptible mice infected with virulent *Brucella abortus* strain 2308. FEMS Immunology & Medical Microbiology 16: 193–203.
 - 9- Klinman DM, Yamshchikov G, Ishigatsubo Y (1997) Contribution of CpG motifs to the immunogenicity of DNA vaccines. J. Immunol. 158:3635–3639.
 - 10- Huang L, Krieg AM, Eller N, Scott DE (1999) Induction and regulation of Th1-inducing cytokines by bacterial DNA, lipopolysaccharide, and heat-inactivated bacteria. Infect. Immun. 67: 6257–6263.
 - 11- Iyer SS, Cheng G (2012) Role of interleukin 10 transcriptional regulation in inflammation and autoimmune disease. Crit Rev Immunol. 32(1):23–63.
 - 12- Fernandes DM, Benson R, Baldwin CL (1995) Lack of a role for natural killer cells in early control of *Brucella abortus* 2308 infections in mice. Infect. Immun. 63:4029-4033.
 - 13- Alton GG, Jones LM, Angus RD, Verger JM. (1988) Techniques for the brucellosis laboratory, INRA, Paris.
 - 14- Solorio Rivera JL, Segura-Correa J.C, Sanchez-Gil LG (2007) Seroprevalence and risk factors for brucellosis of goats in herds of

- Michoacan.Mexico.Prev. Vet.M ed.82,282–290.
- 15- Rahman, AKMA, Saegerman, C, Berkvens, D, Fretin, D, Gani, MO, Ershaduzzaman, M, Ahmed, MU, Emmanuel, A (2013) Bayesian estimation of true prevalence, sensitivity and specificity of indirect ELISA. Rose Bengal Test and Slow Agglutination Test for the diagnosis of brucellosis in sheep and goats in Bangladesh, Preventive Veterinary Medicine 110: pp. 242-252
- 16- Garin-Bastuji B, Blasco JM (1997) Caprine and ovine brucellosis (excluding *B. ovis* infection). In : Manual of standards for diagnostic tests and vaccines, Third edition 1996, OIE, Paris, 350-368.
- 17- Blasco JM, Garin-Bastuji B, Marín CM, Gerbier G, Fanlo J, Jiménez de Bagüés MP, Cau C (1994a) Efficacy of different rose bengal and complement of fixation antigens for the diagnosis of *Brucella melitensis* in sheep and goats. Vet. Rec., 134, 415-420.
- 18- Blasco JM, Marín CM, Jiménez de Bagüés MP, Barberán M, Hernández A, Molina L, Velasco J, Díaz R, Moriyón I (1994b) Evaluation of allergic and serological tests for diagnosis of *Brucella melitensis* in sheep. *J. Clin. Microbiol.*, 32, 1835-1840.
- 19- Salih HMS (2010) Brucellosis in Iraq: Epidemiology, present status, and Challenges in controlling the disease. MSc thesis, College of Veterinary medicine, Kansas State University.
- 20- Al-Talafhah AH, Lafi SQ, Al-Tarazi Y (2003) Epidemiology of ovine brucellosis in Awassi sheep in Northern Jordan, Preventive Veterinary Medicine. 60 (4):297–306.
- 21- Kaoud HA, Zaki MM, El-Dahshan AR, Nasr SA (2010) Epidemiology of Brucellosis Among Farm Animals. Nature and Science, 8 (5):190-197.
- 22- Samadi A, Ababneh MMK., Giadinis N D, Lafi SQ (2010) Ovine and Caprine Brucellosis (*Brucella melitensis*) in Aborted Animals in Jordanian Sheep and Goat Flocks, Veterinary Medicine International, vol. 2010, Article ID 458695, 7 pages, 2010. doi:10.4061/2010/458695.
- 23- Al-Fandi M (2005) Seroepidemiology of caprine Brucellosis in Jordan. Small Ruminant Research, 58(1):13-18.
- 24- Al-Hankawe OKH, Rhaymah MS (2012) Comparison between ELISA and other serological tests for diagnosis of brucellosis in sheep in Ninevah province. *Iraqi J Vet Sci* 26(supplimentII):97-103.
- 25- Corsetti PP, de Almeida LA, Carvalho NB, Azevedo V, Silva TMA, Teixeira HC, *et al* (2013) Lack of endogenous IL-10 enhances production of proinflammatory cytokines and leads to *Brucella abortus* clearance in mice. PLoS One 8(9):e74729.10.1371/journal.pone.0074729.