Kufa Journal for Veterinary Medical Sciences Vol.(7). No.(2) 2016



Kufa Journal for Veterinary Medical Sciences

vetmed@uoKufa.edu. iq



The Effects of Ethanolic Extracts Roots of Zingiber officinale and Leaves of Eruca sativa Extract on reproduction as compare with Tadalafil on Male Rats

Ali A. Al-khayyat * Lubna A. Kafi* Bassam A. A. Al-Shimmary**
*Department of Physiology and Pharmacology, College of Veterinary Medicine,
University of Baghdad.

** Department of Physiology and Pharmacology, College of Veterinary Medicine, University of Kufa.

Abstract

Two ethanolic extracts of *Eruca Sativa* (Rocket) and *Zingiber officinale* (Ginger) were studied after 42 days of treatment with effective dose (ED), as 200 mg per kg of B.W. for rocket and 400 mg per kg of B.W. for ginger, the results were 100% for fertility index(FI) for the two extracts, and, 60% for Tadalafil and 80% for control group. While, the result of pregnancy index(PI)was 100%, that give life babies after successful mating.

Introduction:

All civilizations have always had traditions of using herbs to promote healing. Plants still remain the basis for development of modern drugs and medical plants have been used for years in daily life to treat diseases all over the world [Ates and Erdogrul. 2003]. Ginger has been reported to bind testosterone, raising concerns about possible consequences for the developing embryo(Marcus and Snodgrass, 2005). Also, Amir Amin et al., (2006) that both ethanolic and aqueous extracts of ginger increases serum testosterone level, testicular cholesterol level and activity testicular antioxidant enzymes. Rocket is also considered a medical plant with many reported properties, including its strong aphrodisiac effect known since Roman times [Padulosi, and Pignone, 1997; Font et al., 2003], in addition, Zena, (2013) results exhibited that rocket leaves extract caused a

significant increase in testosterone level, sperm activity; moreover a significant decrease in sperm mortality and abnormalities was recorder.

Materials & methods:

Extract preparation:

Ginger was obtained from commercial source from Najaf city ,and voucher specimen of the plant to be identified and authenticated at the National Herbarium of Iraq Botany directorate Graib-Baghdad Abu under certificate name Zingiber Officinale belongs to family Zingiberaceae, the root powder of Z.O. was put in thimbles that used in multi-units of Soxhlet apparatus to method described by Harborne (1984), with 70% ethanol for 24 hours(Complete circulation) .The combined ethanolic extract was filtered , and the filterate concentrated by a rotary evaporator under a temperature not exceeding 40 C and a speed 80 rpm.

2016

The leaves of Eruca Sativa was purchase from local planter ,that cultivation in Najaf city .The plant classification was done at the Ministry of Agriculture of Iraq/State board of seeds testing and certificate, the powder of ES was prepared by the same procedure which followed with ZO.

Administration of ZOE, ESE and tadalafil:

After 42 days of treatment with ZOE, ESE, and tadalafil dissolved in DMSO as vehicle. The control group was treated with DMSO only, five groups were used for each treatment, each group one male was put with 3 non-treated females, the groups were observed within periodic examination for 3 weeks, until parturition.

Reproductive parameters:

Male rats that were treated with the two ethanolic extracts and tadalafil

were put with untreated females rats mating and the following parameters were considered:

Fertility index = number of pregnancy female rats / number of female mating × 100.

Pregnancy index = number of female that give live fetus / total pregnancy rats \times 100.

Statisvical analysis:

All data were expressed as mean \pm SE . Student's t-test was performed by Statistical Package for the Social Science (SPSS Inc. Chicago, IL,USA) software (version 15.0) with level of significance set at P < 0.05.

Results:

The results of fertility index was 100% for the two ethanolic extracts, however, but 60% for tadalafil as compare with 80% for control group(table 1). While, the results

for pregnancy index was 100% for all groups (table 2) as the following: Table1:Fertility index(FI).

Group	Control	ESE	ZOE	Tadalafil	
No. of female	5	5	5	5	
mating					
successfully					
No. of pregnant	4	5	5	3	
female					
No. of non	1	0	0	2	
pregnant female					
Fertility Index	80%	100%	100 %	60%	
percentage					

Table2:Pregnancy index(PI).

Group	Control	E.S.E.	Z.O.E.	Tadalafil	
No. of pregnant	4	5	5	3	
female					
No. of pregnant	4	5	5	3	
female that gives					
life babies					
Pregnancy Index	100 %	100 %	100%	100 %	
percentage					

Discussion:

The fertility index, showed decrease in percentage in the tadalafil group as compare with the control group, this may be due to decrease in motility of sperm caused by long use of tadalafil, but high percentage in fertility index was observed with the two groups (ZOE and ESE)as compare with the control group, this may be due to good sperm motility ,that results from active ingredients of ginger and rocket on sperm mitochondria.

For ESE this result corroborate with Salem and Moustafa ,(2001) and Zena, (2013), who concluded that Eruca sativa may be capable in improving healthy sperm parameters and fertility. In addition, it is known that ZOE androgenic possess property (Kamtchouing et al, 2002), also, al.,(2000) Kamtchouing et observed an increase in the testicular weight of rats treated with Zingiber Officinale for 8 days with concomitant increase in testosterone level, Another explanation for these improved testicular/sperm function in this study is the reduction in level of malondialdehyde, the most abundant individual aldehyde resulting from peroxidation lipid breakdown biological systems. Malondialdehyde is an indirect indicator of reactive oxygen species, ROS, which has potential toxic effects on sperm quality and function (Sharma and Agarwal, 1996; Sikka et al, 1995).

While, the pregnancy index, showed good ratio in pregnancy index for all groups compare with the control group, that indicate the adverse effects of tadalafil and the active ingredients of the two extracts on female that give life For ZOE, this result is in babies. agreement with Wilkinson, (2000) who stated that recent literatures survey have reported that the most commonly used natural drugs for the treatment of morning sickness is ginger. For ESE in pregnancy index the result is in have agreement with Said et al., (2002)who stated that improvement of antioxidant activity, mitochondrial increasing of and oxidative phosphorylation take place.

References:

	Amir	Amin	, Ala	aa E	Eldin	Α,
Hamz	zaa Aa (2	2006).	Effec	ets O	f Gin	ger
And	Roselle	On	Cispl	atin	Indu	ced
Repro	oductive	Toxic	ity In	Rat	ts. As	sian
J. Āno	drol. 5:60	07 - 6	12.			

- Ates, D., And Erdogrul, O. Antimicrobial Activates Of Various Medicinal And Commercial Plant Turk.J.Biol, Extracts. 27:157-162. 2003.
- Font, R.Galan, S., Ruiz, P., And Villatoro. P. Delrio. Characterization Of The Sensorial, Morphological And Agronomic Attributes Of A World Collection Of Rocket. Brassica, 5thinternational Symposium On Brassica And The 16th Crucifer Genetic Workshop2003.
- Harborne, J.B. (1984).Phytochemical Method: A Guide To Modern Techniques Of Plant Analysis. Champman And Hall, London, Uk.,1-34.
- Kamtchouing P, Fandio Gym, Dimo T, Jatsa Hb. (2002). Evaluation Of Androgenic Activity Of Zingiber Officinale And Pentadiplandra Brazzeana In Male Rats. Asian J Androl. 4:299-301.
- Kamtchouing Ρ, Mbongue Famdio Gy, Dimo T, Jasta Hb (2000). Evaluation Of Androgenic Activity Of Zingiber Officinale Andpentadiplandra Brazzeara In Male. Asian J. Androl. 4: 299 - 301.
- Marcus D, Snodgrass W. Do No Harm: Avoidance Of Herbal Medicines During Pregnancy. Obstet Gynecol. 2005;105:1119-1122.

□ Padulosi, S., Pignone, D.
☐ Padulosi, S., Pignone, D. Rocket: Mediterranean Crop For
Theworld. International Plant Genetic
Resources Institute, Rome, Italy. 1997.
☐ Salem M.A.R. And Moustafa
N.A., Histological And Quantitative
Study Of The Effect Of Eruca Sativa
Seed Oil On The Testis Of Albino Rat,
Egyptian J Hosp Med., 2, 148- 162
(2001).
□ Said O, Khalil K, Fulder S,
Azaizeh H. Ethnobotanical survey of
medicinal herbs of the Middle Eastern
region. J. Ethnopharmacol.
2002;83:251–265.
☐ Sharma Rk, Agarwal A. (1996).
Role Of Reactive Oxygen Species In
Male Infertility. Urology. 48: 835-50.
□ Sikka Sc, Rajasekaran M,
Hellstrom Wjg. (1995). Role Of
Oxidative Stress And Antioxidants In
Male Infertility. J Androl.16: 464-8.
□ Wilkinson JM. What do we
know about herbal morning sickness
treatments? A literature survey.
Midwifery 2000; 16:224–8.
Zena F. Hussein,2013. Study
The Effect Of Eruca Sativa Leaves
Extract On Male Fertility In Albino
Mice. Journal Of Al-Nahrain
University Vol.16 (1), March, 2013,
Pp.143-146.